



Published: October 31, 2023

Citation: Willians L, 2023. Aneurysmal Bone Cyst in the Distal Metaphysis of the Right Fibula: A Case Report , Medical Research Archives, [online] 11(10). <https://doi.org/10.18103/mra.v11i10.4338>

Copyright: © 2023 European Society of Medicine. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

DOI

<https://doi.org/10.18103/mra.v11i10.4338>

ISSN: 2375-1924

CASE REPORT

Aneurysmal Bone Cyst in the Distal Metaphysis of the Right Fibula: A Case Report

Luis Willians MD.

Orthopedic Surgeon – Orthopedic Oncologist.
Associate of the Bone Tumors Service – Padre Machado Oncology Hospital. Caracas, Venezuela

Email: luiswi16@gmail.com

ABSTRACT

21-year-old female patient, she presented increased volume and pain in the right ankle, imaging studies were carried out, evidencing alteration of the bone structure in the distal metaphysis of the fibula, having as a diagnostic option by clinic, epidemiology and imaging, ANEURISMATIC BONE CYST , due to the location of said lesion , and taking into account the percentage of recurrence of the aforementioned lesion , it was decided to perform in bloc resection of the lesion and reconstruction with ipsilateral fibula autograft , descending it and fixing it with osteosynthesis , to maintain the anatomy of the lesion . the ankle joint.

Introduction

The aneurysmal bone cyst (ABC) is a reactive bone lesion characterized by cystic cavities containing blood, often causing light to moderate pain. ABCs can exhibit aggressive growth and cortical destruction, potentially leading to deformity and pathological fractures in affected areas ¹⁻².

The term "aneurysmal bone cyst" (ABC) was first coined by Jaffe and Lichtenstein to describe blood-filled cysts with wide spaces, hemosiderin deposits, giant cells, and some cancellous bone tissue in their walls ³.

ABC accounts for 6% of primary bone lesions and can be associated with benign and malignant conditions, such as giant cell tumors, osteoblastoma, chondroblastoma, chondromyxoid fibroma, fibrous dysplasia, osteosarcoma, fibrosarcoma, or chondrosarcoma ²⁻⁴.

ABC typically occurs in the first two decades of life, representing approximately 1.4% of all primary bone tumors, with an incidence of 0.14 to 0.32 per 100,000 individuals ⁵.

The metaphysis of long bones, including the humerus, femur, tibia, and fibula, is the primary location for ABCs. Patients often present with a painful mass, leading to deformity and functional impairment of the affected limb ⁵⁻⁶.

Treatment options vary but are favorable when detected early. Treatment may involve tumor excision, curettage, and cavity filling with bone graft or cement. In cases with large tumors involving major joints, preserving joint function poses a significant challenge ⁶.

Clinical Case

This case involves a 21-year-old female patient who experienced pain and swelling in the right ankle, starting in March 2021. She sought medical attention in June at the Bone Tumor Services at "Hospital Padre Machado," where X-rays revealed a lithic, expansive injury in the distal metaphysis of the fibula (image 1,2). Laboratory analyses ruled out metabolic conditions.

VACUNAS	1ª Dosis	2ª Dosis	3ª Dosis	Refuerzo	Refuerzo	Refuerzo
B.C.G	05/6/14					
ANTIPOLIOMIELITICA	14/06/14	Regresar 21/4/14	VACUNADO 25/11/14	LOTE	LOTE	LOTE
PENTAVALENTE	21/2/14	Regresar 21/4/14	VACUNADO 25/11/14	LOTE	LOTE	LOTE
ANTIROTAVIRUS	21/2/14	Regresar 21/4/14	VACUNADO	LOTE		
ANTINEUMOCOCO 23 VALENTE	21/2/14	Regresar	VACUNADO	LOTE		
ANTIHEPATITIS B R.N	05/6/14	Regresar 21/4/14	VACUNADO	LOTE		
TRIVALENTE VIRAL (SRP)	8-2-15					
ANTIAMARILICA	8/2/15	LOTE / LABORATORIO				
TOXOIDE TETANICO DIFTERICO		Regresar	VACUNADO	LOTE	LOTE	LOTE
ANTI-INFLUENZA ESTACIONAL	29-01-15	Regresar 3/3/15	VACUNADO	LOTE	LOTE	LOTE
ANTIMENINGOCOCO		Regresar	VACUNADO	LOTE		
OTRAS	Varicela 13/11/14	Hepatitis A 21/4/14	Varicela 21/4/14			
	SPP015470073	416118				

INMUNIZATION SCHEDULE (IMAGE 1)



IMAGE 2

Given the epidemiological, clinical, and imaging data, the patient was diagnosed with an ABC. Due to the high rate of local recurrence associated with less invasive procedures, the surgical treatment chosen was a wide en bloc resection, ipsilateral fibular autograft reconstruction, and fixation with a 2.7 mm distal anatomical plate for the fibula with transdermal fixation.

Surgical Procedure

By July 2021, surgery is performed with the support of the José María Vargas Hospital Osteosynthesis Bank in Caracas. After blood exsanguination of the right lower limb and placement of a tourniquet,

lateral approach to the fibula, the lesion was visualized and confirmed by fluoroscopy, resection was performed with healthy margins of 7 cm in length in the first stage, in the second stage the approach was extended, making a cut to diaphyseal level of the fibula, descending from the fibula to the ankle region, it was fixed with a 2.7 mm 11-hole plate and a transindesmal screw was placed (image 3,4), protecting with a surodigital splint for 10 weeks. Radiological controls were carried out 2 weeks postoperatively. 6 weeks and tenth week, starting partial support at 12 weeks and full load through a guided rehabilitation program at 14 weeks.

Nota: TIBIA IZQUIERDA (1/3 MEDIO)	
BACTERIA 1	
ANTIBIOGRAMA Germen (1)	Staphylococcus aureus
Eritromicina	RESISTENTE
Oxacilina	RESISTENTE
Trimetropin sulfá	SENSIBLE
Ciprofloxacina	SENSIBLE
Gentamicina	SENSIBLE
Levofloxacina	SENSIBLE
Clindamicina	RESISTENTE

IMAGE 3



IMAGE 4



IMAGE 5

The Biopsy report yielded (image 5) **SOLID VARIANT ANEURYSMATIC BONE CYST.**

Patient since December 2021, she resumes her daily activities, with the use of conventional footwear, she

has been clinically and radiologically controlled every 4 months with ap and lateral X-rays, last control in August 2022(image 6).



IMAGE 6



IMAGE 7



IMAGE 8

Discussion

The term "aneurysmal bone cyst" was first introduced by Jaffe and Lichtenstein in 1942, also known as Jaffe-Lichtenstein disease. ABCs can be primary or secondary, often associated with other tumors. Radiography, along with MRI, is effective for diagnosis. ABCs appear as lytic, multilobulated, expansive lesions with trabecular bone, and the diagnosis is usually confirmed histopathologically ⁷⁻⁸⁻⁹.

Although some cases of spontaneous healing have been reported, treatment is often required. Treatment options include surgical curettage with or without bone graft, complete excision, selective arterial embolization, or a combination of these procedures. In this case, en bloc resection and

fibular autograft were chosen to minimize the risk of local recurrence ¹⁰⁻¹¹⁻¹².

Conclusions

Aneurysmal bone cysts are generally benign but can be aggressive and recurrent, requiring multiple surgeries. A comprehensive evaluation, including clinical, epidemiological, and imaging data, is essential to determine the appropriate surgical approach. In this case, wide en bloc resection with fibular autograft reconstruction proved successful, with no signs of recurrence at the latest follow-up in July 2023.

While established protocols exist for managing musculoskeletal oncological injuries, individualized treatment plans should be tailored to each patient's unique circumstances.

References

1. Adam Greenspan and Col. CHAPTER 7 Miscellaneous Tumors and Tumor-like Lesions. Differential Diagnosis in Orthopedic Oncology Edit: 2007 by Lippincott Williams & Wilkins.
2. Schajowicz F. Tumores y Lesiones Seudotumorales de Hueso y Articulaciones. Editorial Médica Panamericana. 1992.
3. Lichtenstein L. Aneurysmal bone cyst. A pathological entity commonly mistaken for giant cell tumor and occasionally for hemangioma and osteogenic sarcoma. 1950.
4. Jaffe HL. Aneurysmal bone cyst. Bull Hosp Joint Dis 1950.
5. Chan G, Arkader A, Kleposki R, Dormans JP. Case report: primary aneurysmal bone cyst of the epiphysis. Clin Orthop Relat Res. 2010.
6. Rapp T, Ward J, Alaia M. Aneurysmal Bone Cyst. Jour Amer Acad of Orthop Surg [revista internet]. 2012 ;20(4):233–241. Disponible en: https://journals.lww.com/jaaos/Abstract/2012/04000/Aneurysmal_Bone_Cyst.10.aspx
7. Mostafa M. Subperiosteal resection of fibular aneurysmal bone cyst. Eur J Orthop Surg Traumatol [revista internet]. 2015; 25 (3): 443 – 50. Disponible en: <https://www.ncbi.nlm.nih.gov/pubmed/25205471>.
8. Ulici A, Nahoi C, Carp A, Fodor I, Dinu C. Surgical Treatment of an Aneurysmal Bone Cyst with Avascular Bone Graft. Chirurgia [revista internet]. 2017; 112 (2): 172-177. Disponible en: <https://www.ncbi.nlm.nih.gov/pubmed/28463677>
9. Zhiqing Zhao, Taiqiang Yan, Wei Guo, Rongli Yang, Xiaodong Tang, Wei Wang. Surgical options and reconstruction strategies for primary bone tumors of distal tibia: A systematic review of complications and functional outcome. J Bone Oncol. [revista en internet]; 14: (100)20-9. Disponible en: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6298939/>
10. Woertler K, Brinkschmidt C. Imaging features of subperiosteal aneurysmal bone cyst. Acta Radiol. 2002.
11. Docquier PL, Glorion C, Delloye C. Quiste óseo aneurismático. EMC - Aparato Locomotor. 2011.