

***Lactococcus lactis* NCDO2118 produces anti-hypertensive GABA and exerts acute hypotensive in spontaneously hypertensive rats**

Authors:

Tessália Diniz Luerce Saraivaa, Gisele Maia Etelvinob, Mariana Oliveirab,
Cassiana Severiano de Sousaa, Marcela Santiago Pacheco de Azevedoa,
Alejandra de Moreno de LeBlanc, Jean Guy LeBlanc, Debmalaya Barhd,
Preetam Ghoshe, Franco Venanzif, Henrique Cesar Pereira Figueiredog,
Robson Augusto Souza Santosb, Vasco Azevedoa*

^a Laboratório de Genética Celular e Molecular, Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais, Belo Horizonte, MG, Brazil

^b Laboratório de Hipertensão, Instituto de Ciências Biológicas, Universidade Federal de Minas Gerais, Belo Horizonte, MG, Brazil

^c Centro de Referencias para Lactobacilos (CERELA-CONICET), Chacabuco 145, T4000ILC San Miguel de Tucuman, Argentina

^d Institute of Integrative Omics and Applied Biotechnology, Nonakuri, Medinipur, WB, India.

^e Department of Computer Science, Virginia Commonwealth University, Richmond, VA, USA.

^f Laboratory of Translational Biology, School of Biosciences and Veterinary Medicine, via Gentile da Varano III, University of Camerino, Italy.

^g Escola de Veterinária, Universidade Federal de Minas Gerais, Belo Horizonte, MG, Brazil.

*** Contact information for**

Corresponding Author:

Vasco Azevedo, Laboratório de
Genética Celular e Molecular
LGCM, Instituto de Ciências
Biológicas, Universidade Federal de
Minas Gerais ICB UFMG, CP 486
CEP 31270 901, Belo Horizonte
MG, Brasil. Tel: 55 31 3409 2610.
vasco@icb.ufmg.br

E- mails:

tdluerce@gmail.com
giseleme@gmail.com
mena_flavia_oliveira@yahoo.com.br
cassissousa@gmail.com
marcela.spachecoazevedo@gmail.com
demoreno@cerela.org.ar
leblancjeanguy@gmail.com
dr.barh@gmail.com
pghosh@vcu.edu
franco.venanzi@unicam.it
figueiredoh@yahoo.com
robsonsant@gmail.com
vascoariston@gmail.com

Abstract

Gamma (γ)-aminobutyric acid (GABA) is a molecule with nutraceutical interest since it is recognized for its health-promoting properties such as hypotensive effects. Some lactic acid bacteria (LAB) have been shown to be able to produce GABA. In the present investigation, *Lactococcus (L.) lactis* NCDO2118, a probiotic LAB, was cultured in a medium supplemented with different concentration of glutamate, the substrate for GABA production. This strain was evaluated in SHRSP (Spontaneously Hypertensive Stroke-Prone Rats) to assess its hypotensive effect. The evaluation showed that GABA production is variable depending on the glutamate concentration and incubation time and that the increased GABA production potentiates its hypotensive effect. This work demonstrates that *L. lactis* NCDO 2118 could be used as a tool for developing health-promoting foods, such as those enriched with GABA as an interesting and novel pharmacological strategy.

Keywords

Lactococcus lactis; Gamma-AminoButyric Acid; GABA; Glutamic Acid Decarboxylase; GAD; Hypertension

1. Introduction

Cardiovascular diseases are responsible for approximately 17 million deaths per year, and more than half of the cases are due to complications of hypertension. High blood pressure, can lead to the damage of the kidneys, eyes, arteries, heart, and brain, thereby affecting cognitive ability and vital biological functions. Changes in diet and lifestyle, as well as adherence to medication, may help in a better control of hypertension (World Health Organization, 2013).

Gamma-aminobutyric acid (GABA) is a non-protein four-carbon amino acid responsible for many beneficial properties on human health, including anti-hypertensive effects. When GABA was injected in the lateral ventricle in an animal model, the arterial pressure and heart rate were decreased (Sasaki et al., 1986). The oral route is also a promising strategy for GABA administration, both in subclinical as well as clinical trials (Elliott

& Hobbiger, 1959; Inoue et al., 2003; Takahashi, Tiba, Iino, & Takayasu, 1955).

Lactic acid bacteria (LAB) are the principal microorganisms that produce GABA (Li & Cao, 2010). This capability is due to the performance of glutamate decarboxylase (GAD, EC 4.1.1.15), an enzyme localized in the cytoplasm that is recruited to the membrane when the pH falls. In the cellular membrane, GAD performs the irreversible alpha-decarboxylation of L-glutamate resulting in GABA production. In this context, GABA can increase ATP production and acid tolerance (Higuchi, Hayashi, & Abe, 1997).

Among the factors that affect the production of GABA by bacteria are temperature, pH, incubation time and different media additives (Gomaa, 2015). The aim of this work was to evaluate different conditions of growth for the bacteria *Lactococcus lactis* NCDO2118, a known GABA-producing LAB (Mazzoli et al., 2010), on GABA production and

evaluate its anti-hypertensive potential. For this latter objective, the bacteria were orally administered in an acute hypertension practical model using SHRSP (Spontaneously Hypertensive Stroke-Prone Rats).

2. Material and Methods

2.1. Bacteria and growth conditions.

L. lactis subsp. *lactis* NCDO 2118 (NCDO) was used in this study. It was activated at 30°C in M17 medium (Difco), containing 0.5% glucose (w/v) (GM17) for 24 hours. Afterwards, bacteria were inoculated 1:10,000 (v/v) in GM17, supplemented or not with glutamate (0.5% and 5% (w/v)) and incubated for 48 hours at 30°C. These cultures were prepared freshly every day during the test period.

2.2. Lactococcus lactis in glutamate culture.

To evaluate the effects of glutamate on *L. lactis*, growth, optical density, pH,

colony-forming unit (CFU/mL) and GABA production were determined at four time points: 0, 6, 24 and 48 hours' incubation at 30°C. Optical density was measured at 580 nm (OD₅₈₀), the pH in digital pH meter, and serial dilutions were performed in GM17 medium and plated in triplicate in solid medium containing 1.5% (w/v) agar to determine CFU/mL after incubation at 30°C for 48 hours. GABA-concentrations in the culture supernatants were determined using the enzymatic GABAase assay as described previously using a commercial kit (Tsukatani, Higuchi, & Matsumoto, 2005).

2.3. Animals

Male SHRSP rats (Masineni, Chander, Singh, Powers, & Stierjr, 2005) 20-24 weeks old (weighing between 245 and 320g) were obtained from Animal House of Hypertension Laboratory of Instituto de Ciências Biológicas da Universidade Federal de Minas Gerais (UFMG). All animals received a balanced

diet *ad libitum*. The experimental groups consisted of a minimum of 5 rats, and the protocols were accepted by the animal protection committee and Ethics Center on Animal Use of UFMG.

2.4. Femoral canulation of spontaneously hypertensive rats

The procedure for catheterization was performed as previously described (Meyer & Fish, 2005). Briefly, the animals were anesthetized intraperitoneally with 2.5% of tribromoethanol, a small incision was performed in the inguinal region of the isolated femoral artery and a polyethylene cannula of 15 cm PE50 to 4 cm PE10 was made and implanted into the abdominal aorta through the femoral artery. After the catheter was secured and exteriorized subcutaneously into the dorse of the animal in the cervical region and the incision was sutured, physiological solution (0.9% NaCl, (w/v)) plus heparin (100 U.I/0.1 mL) was administered. The

cardiovascular parameters of the animals were monitored after 24 hours of surgery.

2.5. Oral treatment and monitoring of cardiovascular parameters in Biopac System

Pulsatile arterial pressure (PAP) was recorded by a signal sent to a transducer connected to the cannula inserted into the abdominal aorta. The pressure fluctuations were amplified and converted into signals captured by a data acquisition system (Biopac System, model MP150) and using the software Acqknowledge v.3.5.7 (Biopac System). Mean arterial pressure (MAP) and heart rate (HR) were recorded every 10 minutes during 6 hours. After the first hour of stabilization of cardiovascular parameters, the animals received 1 mL of single oral administration of LAB by gavage. The monitoring was performed until the sixth hour after treatment administration (Figure 1).




Figure 1. Experimental protocol.

2.6. Monitoring of Blood Pressure

by Radiotelemetry

The animals were anesthetized with tribromoethanol 2.5% (2.5 mg/kg), the catheter-transducer was fixed into the abdominal aorta just above the bifurcation of the iliac arteries, and the sensor was fixed to the abdominal wall. The animals were used in the experiment after 10 days. Arterial pressure was monitored by radio frequency for 30 hours using a telemetry system (Data Sciences International, MN; model TA11-PA C40). The software (Dataquest A.R.T., Gold 2.0) acquired the data to be analyzed.

2.7. Statistical Analysis

Data were analyzed using two-Way ANOVA followed by a Bonferroni posthoc test, and $P < 0.05$ was considered significant. Calculations were performed using the GraphPad Prism 5 software (GraphPad Software, San Diego, CA, USA).

3. Results

3.1. Effect glutamate in *L. lactis* culture

Lactococcus lactis exhibits a different behavior when cultured in GM17 medium supplemented with 0.5% or 5.0 % of glutamate. The optical density, pH and

GABA concentration of supernatant were monitored during 48 hours (Figure 2). The optical density presented larger increase in the 6 hour slot in both culture conditions, while in 24 hours the microorganism were already in the stationary growth phase. No growth was observed in the control

containing only the culture medium. The lowest pH values were observed after the 24 hour and a small increase (of 3 and 7%) was observed in the 48 hour of cultures NCDO in GM17 containing 0.5 and 5.0% glutamate respectively (Figure 2).




Figure 2. Glutamate effect in *Lactococcus* culture.

The production of GABA showed a progressive increase over time. After 24 hours incubation, 30 or 28 mM GABA was produced when *L. lactis* was grown with 0.5 or 5.0 % glutamate respectively.

The highest yield was observed after 48 hours incubation, there was a higher yield with 48 and 65 mM, respectively (Figure 3).




Figure 3. GABA *Lactococcus lactis* NCDO 2118 production.

3.2 Potential hypotensive effect of GABA-producing *Lactococcus lactis*

To evaluate the hypotensor potential effect of *L. lactis* NCDO2118, the bacterial culture was supplemented or not with 5% glutamate. After 48 hours of growth, 1 mL of *L. lactis* was administered orally in SHRSP animals. The arterial pressure was monitored continuously for 6 hours at 10 minutes' intervals. Figure 4 presents the results comparing *L. lactis* groups with (NCDO GGM17) and without (NCDO GM17) glutamate supplementation. The

hypertensive animals treated with only the strain (grown without glutamate) showed a gradual decrease of arterial pressure starting at 60 minutes up to 200 minutes post-administration, where a 22 mmHg decrease was observed. Past this point, a gradual increase of blood pressure was observed until it returned to the initial level. The group with increased production of GABA (NCDO – GGM17), presented a progressive reduction in arterial pressure from the 40th minute with values that were significantly lower than those of the non-glutamate induced group. In this group, the arterial pressure was reduced up

to 28 mmHg after 300 minutes post-inoculation with the microorganisms, and this blood pressure remained constantly lower during the entire evaluation time.

Comparing the effect produced by the two groups on the reduction of hypertension, a statistical difference was found between treatments.




Figure 4. Graphical depiction of Mean Arterial Pressure (MAP) with *Lactococcus lactis* NCDO2218 treatment in SHRSP rats. NCDO GM17- hypertensive group, treated with *L. lactis* without glutamate supplementation; and NCDO GGM17- hypertensive group treated with *L. lactis* with glutamate supplementation. *** P<0.01

When administering either the supernatant or cell fractions in hypertensive animals, none of these fractions alone was able to produce the same effect that was observed when the

whole culture of *L. lactis* grown with 5% glutamate was used. The treatment with entire culture was statistically more effective against hypertension than any of the isolated fractions (Figure 5).




Figure 5. Graphical depiction of Mean Arterial Pressure (MAP) with *Lactococcus lactis* NCDO2218 culture and its fractions in SHRSP rats treatment. NCDO GGM17- hypertensive group, treated with *L. lactis* with glutamate supplementation; NCDO sup - hypertensive group treated with supernatant of *L. lactis* culture with glutamate supplementation; and NCDO cell - hypertensive group treated with cell fraction of *L. lactis* culture with glutamate supplementation.

*** P<0.01

3.3. Monitoring of blood pressure reduction by telemetry

To evaluate the effect of *L. lactis* strain on elevated blood pressure in rats for more than 6 hours, a telemetry system

was used and was monitored for 30 hours. During this period, the animals of all groups showed an increase in blood pressure from the tenth hour followed by a slight reduction (Figure 6).




Figure 6. Graphical depiction of telemetry measurements to effect of the treatment with *Lactococcus lactis* NCDO2218 for 30 hours. GGM17 - hypertensive group treated with culture medium with glutamate supplementation; NCDO GM17- hypertensive group treated with *L. lactis* without glutamate supplementation; and NCDO GGM17- hypertensive group treated with *L. lactis* with glutamate supplementation.

4. Discussion

Gamma-Aminobutyric acid (GABA) is an amino acid that acts as an inhibitory neurotransmitter in the nervous system. The GABA system regulates important psychological processes, including emotional behavior (Bravo et al., 2011); dysfunction of such processes may involve anxiety and depression (Fatemi, Folsom,

Rooney, & Thuras, 2013). GABA also regulates, among other functions, cardiovascular conditions such as blood pressure and heart rate (Cryan & Kaupmann, 2005; Mody, De Koninck, Otis, & Soltesz, 1994; Roberts & Frankel, 1950; Schousboe & Waagepetersen, 2007).

The ability of GABA in lowering blood pressure has been reported in animal and human trials. Its potential is attributed in part to blocking the peripheral ganglia, or possibly through the inhibition of noradrenaline release from sympathetic nerve endings (Elliott & Hobbiger, 1959; Hayakawa, Kimura, & Kamata, 2002; Stanton, 1963; Takahashi et al., 1955).

An interesting pharmacological strategy is the modulation of GABAergic activity through a system involving a rapid uptake of GABA into presynaptic terminals and surrounding glial cells (Krogsgaard-Larsen, Falch, Larsson, & Schousboe, 1987; Thwaites, Basterfield, McCleave, Carter, & Simmons, 2000).

GABA is naturally present in many foods such as vegetables and fruits (Takayama & Ezura, 2015), especially in fermented products. Since GABA is considered a health-promoting functional compound, many dietary supplements and enriched foods have been developed to

increase GABA intake. (Kook & Cho, 2013).

In Europe, the addition of GABA is considered a "dietary supplement" while in the United States it is considered a "food constituent". In Japan (Sanders, 1998), foods fortified with GABA can be denominated "foods for specified health use" (FOSHU), which have received great interest due to the increasing amounts of cases of hypertension in the general population, that is caused at least partially by the consumption of excessive amounts of sodium. Thus, many products have been developed to serve as vehicles for delivery of GABA (Becerra-Tomás et al., 2015). The enrichment of foods are based essentially on fermentation of substrates with GABA-producing bacteria (Inoue et al., 2003; Li, Qiu, Huang, & Cao, 2010). In this regard, dairy, soybean, kimchi and juice products could be used as potential GABA-delivery vehicles (Chang et al., 2009; Kim, Lee, Ji, Lee, & Hwang, 2009; Park & Oh, 2007; Seok et al., 2008). A

fermented milk product containing GABA was shown to have a blood pressure lowering effect in spontaneously hypertensive rats (Hayakawa et al., 2004) and in mildly hypertensive people (Inoue et al. 2003).

Lactococcus lactis, i.e., the LAB model, has scantily been explored for this purpose. Thus, we chose a probiotic strain of *L. lactis* (NCDO2118 or NCDO) to optimize GABA production. Two concentrations of glutamate supplementation (0.5% and 5.0%) were evaluated. The bacterial culture with the highest concentration of glutamate showed a higher optical density in addition to the increase of GABA production over time; this increased growth could be attributed to the fact that glutamate can contribute favorably to the cell power generation. The alpha-ketoglutarate and succinate are intermediates of the Krebs cycle and hence related to cellular energy production. Glutamate can be deaminated to produce alpha-ketoglutarate. Alternatively, via

"GABA shunt" glutamate may be decarboxylated to produce GABA; for this, succinic semialdehyde is converted to succinate, and is donated to an amino moiety that promotes alpha-ketoglutarate glutamate production. Alternatively, energy can be produced from glutamate since the enzymes are dependent on pyridoxal phosphate (Feehily & Karatzas, 2013; Waagepetersen, Sonnewald, & Schousboe, 2002).

After internalization of glutamate from the cell and GAD activity, the intracellular GABA is produced from the excess proton consumption in the midst of high-stress acid in the cytoplasm. The action of GAD is related to acid resistance in order to maintain cytoplasmic pH constant. A possible explanation for the observed slight increase in the pH at 48 hours, may be due to the maintenance of pH by GAD since the highest concentration of GABA production was observed at this time. Moreover in bacteria, GABA can be exported by the

presence of two glutamate\GABA Antiporters, contributing to cellular homeostasis.

After showing that the addition of glutamate significantly increased GABA production, the beneficial effect of the probiotic bacteria was evaluated in acute hypertension using a SHRSP rodent model hypertension.

L. lactis NCDO 2118 grown without glutamate is able to produce low concentrations of GABA. The hypertensive animals treated with this strain showed a progressive reduction in blood pressure with a maximum effect 2.5 hours after the start of pressure reduction. After this time, it took over approximately two more hours to restore hypertensive patterns of the animals tested. On the other hand, when the animals were treated with the same strain grown in high concentrations of glutamate that induced higher concentrations of GABA, a more pronounced effect on decreasing blood pressure was observed. Unlike what

happened with the previous group, the reduction in mean arterial pressure was maintained until the final hours of the experiment. Although the results were favorable in this experiment while evaluating the beneficial effect of GABA in acute hypertension using the Biopac system, they were not observed when assessed using another analysis system. This conflicting data could be explained by the difference of recovery periods of the animal after anesthesia. Using the Biopac system for recording blood pressure values, the rats were submitted to the experimental treatment 24 hours after cannulation surgery, and in telemetry the experiments started 10 days after surgery, because telemetry surgery is a very complex procedure and animals requires a longer recuperation period. As TBE is an anesthetic that exerts sedative and hypnotic effects through GABA system, a short period after anesthesia the GABA receptors could be more sensitive to

GABA (Krasowski & Harrison, 2000; Sauguet et al., 2013).

The mechanism by which GABA can exert its effect is unclear. Oral administration of GABA may be promoting any direct interaction in the brain, or their action is mediated through indirect ways such as by the enteric nervous system (ENS) (Boonstra et al., 2015). The ENS comprises a complex neural network located in the wall of the intestine, which is able to regulate intestine functions independently of the CNS (Auteri, Zizzo, & Serio, 2015). It is also known that following ingestion, GABA transepithelial transport across the intestinal wall can be mediated via a pH-dependent mechanism (H^+ / GABA symport) (Thwaites et al., 2000). The GABA receptor is distributed by ENS, both in the submucosa as neurons and in myenteric (Poulter, Singhal, Brown, &

Krantis, 1999). Then, the increase of GABA level in the blood flow can explain the reduction effect on blood pressure observed in hypertensive rats.

5. Conclusion

In this work, the importance of the GABA produced by a probiotic LAB in the control of hypertension was suggested, showing that GABA has the potential to be used as a nutraceutical. However, more studies are needed to determine the GABA inflow tract, responsible for its beneficial effect. The optimization of probiotic bacteria producing GABA, is an interesting target for the development of foods enriched for therapeutic purposes.

Acknowledgements

The authors declare that they have no competing interests.

References

- Auteri, M., Zizzo, M. G., & Serio, R. (2015). GABA and GABA receptors in the gastrointestinal tract: from motility to inflammation. *Pharmacological Research*, 93, 11–21.
<http://doi.org/10.1016/j.phrs.2014.12.001>
- Becerra-Tomás, N., Guasch-Ferré, M., Quilez, J., Merino, J., Ferré, R., Díaz-López, A., ... Salas-Salvadó, J. (2015). Effect of Functional Bread Rich in Potassium, γ -Aminobutyric Acid and Angiotensin-Converting Enzyme Inhibitors on Blood Pressure, Glucose Metabolism and Endothelial Function: A Double-blind Randomized Crossover Clinical Trial. *Medicine*, 94(46), e1807.
<http://doi.org/10.1097/MD.00000000000001807>
- Boonstra, E., de Kleijn, R., Colzato, L. S., Alkemade, A., Forstmann, B. U., & Nieuwenhuis, S. (2015). Neurotransmitters as food supplements: the effects of GABA on brain and behavior. *Frontiers in Psychology*, 6.
<http://doi.org/10.3389/fpsyg.2015.01520>
- Bravo, J. A., Forsythe, P., Chew, M. V., Escaravage, E., Savignac, H. M., Dinan, T. G., ... Cryan, J. F. (2011). Ingestion of Lactobacillus strain regulates emotional behavior and central GABA receptor expression in a mouse via the vagus nerve. *Proceedings of the National Academy of Sciences*, 108(38), 16050–16055.
<http://doi.org/10.1073/pnas.1102999108>
- Chang, C.-T., Hsu, C.-K., Chou, S.-T., Chen, Y.-C., Huang, F.-S., & Chung, Y.-C. (2009). Effect of fermentation time on the antioxidant activities of tempeh prepared from fermented soybean using *Rhizopus oligosporus*. *International Journal of Food Science & Technology*, 44(4), 799–806. <http://doi.org/10.1111/j.1365-2621.2009.01907.x>
- Cryan, J., & Kaupmann, K. (2005). Don't worry "B" happy!: a role for GABA(B) receptors in anxiety and

- depression. *Trends in Microbiology*, 114(1), 11–24.
<http://doi.org/10.1111/j.1365-2672.2012.05434.x>
- Dhakal, R., Bajpai, V. K., & Baek, K.-H. (2012). Production of gaba (γ - aminobutyric acid) by microorganisms: a review. *Brazilian Journal of Microbiology*, 43(4), 1230–1241.
<http://doi.org/10.1590/S1517-83822012000400001>
- Elliott, K., & Hobbiger, F. (1959). Gamma aminobutyric acid: circulatory and respiratory effects in different species: re-investigation of the anti-strychnine action in mice. *The Journal of Physiology*, 146, 74–80.
- Fatemi, S. H., Folsom, T. D., Rooney, R. J., & Thuras, P. D. (2013). Expression of GABAA α 2-, β 1- and ϵ -receptors are altered significantly in the lateral cerebellum of subjects with schizophrenia, major depression and bipolar disorder. *Translational Psychiatry*, 3(9), e303.
<http://doi.org/10.1038/tp.2013.64>
- Feehily, C., & Karatzas, K. A. G. (2013). Role of glutamate metabolism in bacterial responses towards acid and other stresses. *Journal of Applied Microbiology*, 114(1), 11–24.
<http://doi.org/10.1111/j.1365-2672.2012.05434.x>
- Gomaa, E. Z. (2015). Enhancement of γ -Aminobutyric Acid Production by Co-Culturing of Two Lactobacilli Strains. *Asian Journal of Biotechnology*, 7(3), 108–118.
<http://doi.org/10.3923/ajbkr.2015.108.118>
- Hayakawa, K., Kimura, M., & Kamata, K. (2002). Mechanism underlying γ -aminobutyric acid-induced antihypertensive effect in spontaneously hypertensive rats. *European Journal of Pharmacology*, 438(1-2), 107–113.
[http://doi.org/10.1016/S0014-2999\(02\)01294-3](http://doi.org/10.1016/S0014-2999(02)01294-3)
- Hayakawa, K., Kimura, M., Kasaha, K., Matsumoto, K., Sansawa, H., & Yamori, Y. (2004). Effect of a γ -aminobutyric acid-enriched dairy product on the blood pressure of spontaneously hypertensive and normotensive Wistar–Kyoto rats. *British Journal of Nutrition*, 92(03), 411.
<http://doi.org/10.1079/BJN20041221>

- Higuchi, T., Hayashi, H., & Abe, K. (1997). Exchange of glutamate and gamma-aminobutyrate in a Lactobacillus strain. *Journal of Bacteriology*, 179(10), 3362–3364.
<http://doi.org/10.5851/kosfa.2013.33.3.377>
- Inoue, K., Shirai, T., Ochiai, H., Kasao, M., Hayakawa, K., Kimura, M., & Sansawa, H. (2003). Blood-pressure-lowering effect of a novel fermented milk containing γ -aminobutyric acid (GABA) in mild hypertensives. *European Journal of Clinical Nutrition*, 57(3), 490–495.
<http://doi.org/10.1038/sj.ejcn.1601555>
- Kim, J. Y., Lee, M. Y., Ji, G. E., Lee, Y. S., & Hwang, K. T. (2009). Production of γ -aminobutyric acid in black raspberry juice during fermentation by Lactobacillus brevis GABA100. *International Journal of Food Microbiology*, 130(1), 12–16.
<http://doi.org/10.1016/j.ijfoodmicro.2008.12.028>
- Kook, M.-C., & Cho, S.-C. (2013). Production of GABA (gamma amino butyric acid) by Lactic Acid Bacteria. *Korean Journal for Food Science of Animal Resources*, 33(3), 377–389.
<http://doi.org/10.5851/kosfa.2013.33.3.377>
- Krasowski, M. D., & Harrison, N. L. (2000). The actions of ether, alcohol and alkane general anaesthetics on GABA_A and glycine receptors and the effects of TM2 and TM3 mutations. *British Journal of Pharmacology*, 129(4), 731–743.
<http://doi.org/10.1038/sj.bjp.0703087>
- Krogsgaard-Larsen, P., Falch, E., Larsson, O., & Schousboe, A. (1987). GABA uptake inhibitors: relevance to antiepileptic drug research. *Epilepsy Research*, 1(2), 77–93.
- Li, H., & Cao, Y. (2010). Lactic acid bacterial cell factories for gamma-aminobutyric acid. *Amino Acids*, 39(5), 1107–1116.
<http://doi.org/10.1007/s00726-010-0582-7>
- Li, H., Qiu, T., Huang, G., & Cao, Y. (2010). Production of gamma-aminobutyric acid by Lactobacillus brevis NCL912 using fed-batch fermentation. *Microbial Cell Factories*, 9(1), 85.
<http://doi.org/10.1186/1475-2859-9-85>

- Masineni, S., Chander, P., Singh, G., Powers, C., & Stierjr, C. (2005). Male Gender and Not the Severity of Hypertension Is Associated With End-Organ Damage in Aged Stroke-Prone Spontaneously Hypertensive Rats. *American Journal of Hypertension*, 18(6), 878–884. <http://doi.org/10.1016/j.amjhyper.2004.12.014>
- Mazzoli, R., Pessione, E., Dufour, M., Laroute, V., Giuffrida, M. G., Giunta, C., ... Loubière, P. (2010). Glutamate-induced metabolic changes in *Lactococcus lactis* NCDO 2118 during GABA production: combined transcriptomic and proteomic analysis. *Amino Acids*, 39(3), 727–737. <http://doi.org/10.1007/s00726-010-0507-5>
- Meyer, R., & Fish, R. (2005). A review of tribromoethanol anesthesia for production of genetically engineered mice and rats. *Lab Animal*, 34(10), 47–52.
- Mody, I., De Koninck, Y., Otis, T. S., & Soltesz, I. (1994). Bridging the cleft at GABA synapses in the brain. *Trends in Neurosciences*, 17(12), 517–525. [http://doi.org/10.1016/0166-2236\(94\)90155-4](http://doi.org/10.1016/0166-2236(94)90155-4)
- Park, K.-B., & Oh, S.-H. (2007). Production of yogurt with enhanced levels of gamma-aminobutyric acid and valuable nutrients using lactic acid bacteria and germinated soybean extract. *Bioresource Technology*, 98(8), 1675–1679. <http://doi.org/10.1016/j.biortech.2006.06.006>
- Poulter, M. O., Singhal, R., Brown, L. A., & Krantis, A. (1999). GABAA receptor subunit messenger RNA expression in the enteric nervous system of the rat: implications for functional diversity of enteric GABAA receptors. *Neuroscience*, 93(3), 1159–1165. [http://doi.org/10.1016/S0306-4522\(99\)00174-8](http://doi.org/10.1016/S0306-4522(99)00174-8)
- Roberts, E., & Frankel, S. (1950). γ -Aminobutyric acid in brain. *Federation Proceedings*, (9), 219.
- Sanders, M. E. (1998). Overview of Functional Foods: Emphasis on Probiotic Bacteria. *International Dairy Journal*, 8(5-6), 341–347.

- [http://doi.org/10.1016/S0958-6946\(98\)00056-9](http://doi.org/10.1016/S0958-6946(98)00056-9)
- Sasaki, H., Muramoto, O., Kanazawa, I., Arai, H., Kosaka, K., & Iizuka, R. (1986). Regional distribution of amino acid transmitters in postmortem brains of presenile and senile dementia of Alzheimer type. *Annals of Neurology*, 19(3), 263–269.
- Sauguet, L., Howard, R. J., Malherbe, L., Lee, U. S., Corringer, P.-J., Adron Harris, R., & Delarue, M. (2013). Structural basis for potentiation by alcohols and anaesthetics in a ligand-gated ion channel. *Nature Communications*, 4, 1697. <http://doi.org/10.1038/ncomms2682>
- Schousboe, A., & Waagepetersen, H. S. (2007). GABA: Homeostatic and pharmacological aspects. In *Progress in Brain Research* (Vol. 160, pp. 9–19). Elsevier. Retrieved from <http://linkinghub.elsevier.com/retrieve/pii/S0079612306600022>
- Seok, J., Park, K., Kim, Y., Bae, M., Lee, M., & Oh, S. (2008). Production and Characterization of Kimchi with Enhanced Levels of γ -Aminobutyric Acid, 17(5).
- Stanton, H. (1963). Mode of action of gamma amino butyric acid on the cardiovascular system. *Archives Internationales de Pharmacodynamie et de Thérapie*, 143, 195–204.
- Takahashi, H., Tiba, M., Iino, M., & Takayasu, T. (1955). The effect of γ -aminobutyric acid on blood pressure. *The Japanese Journal of Physiology*, 5, 334–341.
- Takayama, M., & Ezura, H. (2015). How and why does tomato accumulate a large amount of GABA in the fruit? *Frontiers in Plant Science*, 6. <http://doi.org/10.3389/fpls.2015.00612>
- Thwaites, D. T., Basterfield, L., McCleave, P. M. J., Carter, S. M., & Simmons, N. L. (2000). Gamma-aminobutyric acid (GABA) transport across human intestinal epithelial (Caco-2) cell monolayers. *British Journal of Pharmacology*, 129(3), 457–464. <http://doi.org/10.1038/sj.bjp.0703069>
- Tsukatani, T., Higuchi, T., & Matsumoto, k. (2005). Enzyme-based microtiter plate assay for γ -aminobutyric acid: application to the screening of γ -

- aminobutyric acid-producing lactic acid bacteria. *Analytica Chimica Acta*, 540, 293–297.
- Waagepetersen, H. S., Sonnewald, U., & Schousboe, A. (2002). The GABA Paradox: Multiple Roles as Metabolite, Neurotransmitter, and Neurodifferentiative Agent. *Journal of Neurochemistry*, 73(4), 1335–1342. <http://doi.org/10.1046/j.1471-4159.1999.0731335.x>
- World Health Organization. (2013). *A Global Brief on Hypertension: Silent Killer, Global Public Health Crisis* (online). Retrieved from <http://www.thehealthwell.info/node/466541>